

Performance Recording – “Getting it Right”

The term “rubbish in – rubbish out” is often used when discussing the BREEDPLAN analysis. In other words, the reliability of the EBVs that are produced is a direct reflection of the reliability (or quality) of the performance information that is submitted to BREEDPLAN.

This document outlines 10 of the most common issues that cause seedstock producers problems when performance recording with BREEDPLAN. Importantly, the steps that seedstock producers can take to avoid these problems are also detailed. Careful consideration of this information allows producers to take a large step towards ensuring that they are maximising the returns from their investment in performance recording.

Problem 1 – Inadequate Planning

The cause of a considerable number of performance recording problems stem back to inadequate planning (or inadequate priority) being given to the performance recording requirements of a seedstock herd.

In this scenario, little planning goes into the management of the herd so that the value of any collected performance data is maximised. Herds will often not record the required performance, will record performance information in an ad hoc manner and in a lot of situations, performance information will only be collected following a letter from BREEDPLAN outlining submission deadlines for inclusion in GROUP BREEDPLAN analyses.

In layman’s terms, these herds may be classed as “reactive” or “casual” performance recorders.

Solutions

- ✓ Become a “proactive” performance recorder
- ✓ Incorporate the collection of performance information into the standard management of your herd.
- ✓ Plan data collection ahead of time
- ✓ Submit data to BREEDPLAN shortly after data collection

Problem 2 – Poor Recording of Recipient Dam Information

There are now a considerable number of seedstock producers using embryo transfer within their breeding program. While BREEDPLAN has the ability to analyse the performance of embryo transfer calves, inadequate recording of recipient dam information or use of recipient dams of different breeds often results in performance recording problems.

Importantly, the amount of information available on the recipient dam determines how BREEDPLAN uses the performance information of each ET calf.

Solutions

- ✓ Use recipient dams that are all the same breed when running an ET program. (BREEDPLAN will only directly compare the performance of ET calves that have been reared by recipient dams of the same breed)
- ✓ Ensure adequate recipient dam information is recorded with your Breed Society/Association for each ET calf (ie. tag no., specific breed and year of birth)
- ✓ Ideally, use “known” recipient dams. This includes recipient dams that have been used in previous ET programs or alternatively, stud cows that have Milk EBVs available.

Problem 3 – Unverified Outliers

As part of the on going verification of the performance information that is collected, BREEDPLAN checks the variation in performance records between animals within each contemporary group. While a certain degree of variation is expected within each group, when the difference between a performance record for an animal and the average of all animals in that contemporary group is greater than expected, the record for the animal is flagged as an outlier.

Each time an “outlier” is identified, an outlier report is forwarded to the relevant herd. This report gives the breeder the opportunity to correct or verify the performance for the “outlier” animal. If BREEDPLAN receives no response to the outlier report, the outlier records are excluded from all future GROUP BREEDPLAN analyses.

Ignoring outlier reports is a common cause of performance recording problems.

Solutions

- ✓ Pay attention to all information that you receive from the BREEDPLAN office.
- ✓ Verify/Correct all outlier reports as soon as you receive them in the mail. If you are in any doubt, contact your BREEDPLAN processor at ABRI for advice.

Problem 4 – Inadequate Recording of Important Traits

BREEDPLAN currently has the potential to produce up to 18 different EBVs on each particular animal. While an EBV must meet minimum accuracy criteria (and therefore, be of adequate reliability) before it will report, EBVs for some traits may appear for individual animals when minimal direct performance data recorded has been recorded for that trait.

Common examples include:

- No birth weight has been recorded but birth weight EBVs are reporting.
- Only a 200 day weight has been recorded but all weight EBVs are reporting.
- No 200 day weights have been recorded in a herd but Milk EBVs are reporting on all dams and sires.
- Only weight traits have been recorded but carcass EBVs (e.g. IMF%, EMA) are reporting.

While it is possible to generate “reliable” EBVs from pedigree information and performance that has been recorded for correlated traits, generally speaking EBVs will be of lower reliability if animals haven’t been directly recorded for the trait.

Problems often occur when seedstock producers don’t adequately record the traits of importance to them and their clients.

Solutions

- ✓ Record all traits that are important to either you or your clients
- ✓ Do not rely on pedigree and minimal trait recording to generate EBVs.

Problem 5 – Poor Genetic Linkage

Genetic linkage across contemporary groups both within a herd and between different herds is of fundamental importance in allowing the generation of GROUP BREEDPLAN EBVs. Unfortunately, some common management practices can reduce genetic linkage significantly. More specifically, such management practices include:

(i) Within herd

- Completely replacing all sires from one joining season to the next
- Managing the calves from one sire differently to all other calves before recording any performance information
- Not mixing cows after joining
- All ET calves being by the same sire (and dam)

(ii) Across herd

- All sires used in the herd having no performance recorded progeny in any other herds (for a range of traits).
- Managing the calves from one sire (AI) differently to others before recording traits

- AI calves in a herd being born at a separate time to those calves from natural matings.

Poor genetic linkage can cause significant performance recording problems.

Solutions

- ✓ Do not replace all sires from one year to the next so that across year comparisons can be made.
- ✓ Use sires that have progeny recorded for a range of traits in other herds.
- ✓ Mix cows after joining, particularly AI females.
- ✓ Manage calves by a range of sires together until after the key performance traits have been recorded
- ✓ In ET programs, try to have a range of sires represented
- ✓ Ensure calves from an AI program are born at a similar time to naturally conceived calves by your home bred sires

Problem 6 – Small Contemporary Groups

Although the BREEDPLAN analysis is a very complex analytical model, the basic mechanism by which it works is to directly compare the performance of an animal with the performance of other “similar” animals within the same contemporary group.

Where only a small number of animals are represented in a contemporary group, there are only a few “similar” animals to which it’s performance can be directly compared and thus the performance submitted for it can not be used effectively by the BREEDPLAN analysis.

Small contemporary groups are a problem frequently experienced by smaller herds and without careful management, can result in considerable performance recording problems.

Solutions

- ✓ Restrict calving periods. A calving period of 6 to 8 weeks is optimal.
- ✓ Run as many calves as possible under the same management conditions.
- ✓ Weigh all animals in a management group on the same day.
- ✓ If a management group has to be split, weigh all calves before splitting the group eg. all males before they are castrated.

- ✓ Create management groups based on “automatic” criteria eg sex, prior management groups, prior weigh dates.
- ✓ If you have a commercial herd of similar breed content to your stud animals, it may also be possible to record these animals with your relevant Breed Association/Society.
- ✓ In the situation where two herds run their animals together on the same property, set up an associate membership with BREEDPLAN.

Problem 7 – Single Sire Contemporary Groups

In the same way that it is important to have more than one calf represented in each contemporary group, it is also important to have the progeny from more than one sire represented within a contemporary group.

Where all calves in a contemporary group are by the same sire, there are no other calves by other sires to which the performance of these calves can be directly compared. In this manner, the performance submitted for those calves can not be used effectively by the BREEDPLAN analysis to calculate the EBVs of their sire.

Single sire contemporary groups are a problem that can be experienced by herds of all sizes and without careful management, can result in considerable performance recording problems.

Solutions

- ✓ Use more than one sire in each particular joining.
- ✓ Mix cows after joining
- ✓ Consciously manage the herd so that more than one sire is represented in each contemporary group.

Problem 8 – Selective Performance Recording

Significant problems also arise when only a selection of animals in a contemporary group are performance recorded. For example, only the performance for the best animals in the contemporary group is submitted to BREEDPLAN. In this situation, the performance information for an animal will only be compared with the “selection” that has been recorded. If this “selection” is not an accurate reflection of the entire contemporary group, then BREEDPLAN can not make adequate comparisons and the EBVs produced may be biased or misleading.

The table on the following page provides an illustration of the problems caused by selective recording.

Animal	200 Day Wt (1)	200 Day Wt (2)
A1	255	255
A2	238	238
A3	261	261
A4	205	X
A5	187	X
A6	265	265
A7	237	237
A8	195	X
A9	258	258
A10	228	X
Average	233 kg	252 kg

The problem caused by selective recording is demonstrated if we consider animal A7. In the first scenario, all 10 calves in the contemporary group have been recorded and A7 is 4 kg heavier than the average of the group (237 kg v's 233 kg). This is an accurate reflection of how this animal ranks compared to his peers.

However, in the second scenario, the weights for the lightest 4 calves have not been recorded. A7 is now 15 kg lighter than the "average" of the group (237 kg v's 252 kg). The selective recording of this group has resulted in the performance of A7 being compared against a misleading average. As BREEDPLAN can only calculate EBVs based on the information that has been recorded, the subsequent EBVs that are calculated from this performance will also be misleading (ie. biased).

Solutions

- ✓ Adopt a whole herd recording strategy
- ✓ Record all calves with your Breed Society/Association
- ✓ Always record performance for all calves in a contemporary group and submit this performance to BREEDPLAN
- ✓ Record performance on all available animals (ie. heifers, bulls & steers, rather than just bulls).

Problem 9 – Over Management Grouping

As identified in "Problem 6", one common problem encountered is the separation of calves into isolated contemporary groups of only one or two animals (thereby virtually eliminating those calves from any comparison with their peers). While this is a problem in itself when calves are managed in small groups, issues can also arise when calves are part of a large contemporary group but their performance is analysed in small contemporary groups by BREEDPLAN.

The common cause of this problem is the submission of many different management groups - even though calves are being run as part of the one large contemporary group.

Solutions

- ✓ Understand the criteria that automatically form contemporary groups in BREEDPLAN (eg. herd, calving year, sex)
- ✓ Only submit management groups for non-genetic factors (eg. sick animals, animals run under different conditions).

Problem 10 – Inadequate Management Grouping

BREEDPLAN analyses cattle in contemporary groups to take out the influence of as many of the non-genetic effects as possible (eg. feeding, years, seasons). The underlying principle is that only animals that have had an equal opportunity to perform are directly compared together within each contemporary group.

If the contemporary groups are not correctly formed, the EBVs calculated will be less accurate and possibly misleading. Most of the problems that breeders encounter in “believing” their BREEDPLAN EBVs can be traced back to incorrect management grouping. Poor management grouping will result in BREEDPLAN not being able to differentiate between calves that have had different levels of management or feeding.

Solutions

- ✓ Understand the importance and role of management groups
- ✓ Provide management groups for those animals that have been treated differently to their peers and have performed differently due to the non-genetic factors.
- ✓ If you are in any doubt, contact staff at BREEDPLAN for advice

For more information regarding any of the performance recording problems discussed above, please contact staff at BREEDPLAN